

**BERESKIN & PARR**

**UNITED STATES APPLICATION**

**Title: IMMUNOGENIC PEPTIDES DERIVED FROM PROSTATE-SPECIFIC ANTIGEN (PSA) AND USES THEREOF**

**Inventors: Artur Pedyczak, Pele Chong, Charles D Y Sia**

**Title: IMMUNOGENIC PEPTIDES DERIVED FROM PROSTATE-SPECIFIC ANTIGEN (PSA) AND USES THEREOF.**

The application claims benefit from U.S. Provisional Application Serial No. 60/195,456 filed April 10, 2000 which is incorporated herein by reference in its entirety.

**FIELD OF THE INVENTION**

The present invention relates to immunology, in particular to peptides derived from prostate-specific antigen (PSA) and nucleic acids coding therefor, recombinant nucleic acids into which are inserted nucleic acids coding for PSA peptides, and their use as immunogenic agents.

**BACKGROUND OF THE INVENTION**

Cancer of the prostate is the most commonly diagnosed cancer in man and is the second most common cause of cancer death (Carter, H.B. and Coffey, D.S. (1990) *Prostate* 16:39-48; Armbruster, D.A. (1993) *Clin Chem* 39:181-195). If detected at an early stage, prostate cancer is potentially curable. However, a majority of cases are diagnosed at later stages when metastasis of the primary tumor has already occurred (Wang, M.C., Kuriyama, M., Papsidero, L.D., Loo, R.M., Valenzuela, L.A., Murphy, G.P., and Chu, T.M. (1982) *Methods in Cancer Research* 19:179-197). Present treatments for prostate cancer include radical prostatectomy, radiation therapy, or hormonal therapy. No systemic therapy has clearly improved survival in cases of hormone refractory disease. With surgical intervention, complete eradication of the tumor is not always achieved and the observed reoccurrence of the cancer (12-68%) is dependent upon the initial clinical tumor stage (Zietman, A.L., Shipley, W.L., and Willett, C.G. (1993) *Cancer* 71:959-969). Thus, alternative methods of treatment including prophylaxis or prevention are desirable.

The advent of DNA technology and its expanding use in the field of immunology has led to the identification of human tumor-associated antigens (TAAs) (Rosenberg, S.A. (1995) *Cancer J. Sci. Am.* 12:89-100; Boon, T. (1993) *Int. J. Cancer* 54:177-180). These TAAs now offer the potential of applying modern immunotherapeutic approaches to the treatment of some

human cancers. The basis of one such approach is to immunize a host with appropriate immunogen(s) to elicit cell-mediated immunological responses involving recruitment of tumor-specific effector cells (e.g. cytotoxic T lymphocytes (CTLs)) in an effort to recognize and destroy neoplastic cells.

5       The human TAAs that have been identified to date can be classified into four general categories. The first encompasses the 'cancer/testis' antigens, such as those of the MAGE gene family, whose expression is tumor-specific. The second encompasses antigens which are virally-derived, such as those from human papilloma virus (HPV) and Epstein-Barr virus  
10 (EBV). The third encompasses differentiation antigens, including the prostate-specific antigen (PSA), prostate-specific membrane antigen (PSMA), Melan-A/MART-1, tyrosinase, gp100 and ganglioside GM2. The fourth encompasses antigens such as modified/mutated  $\beta$ -catenin, ras and p53, which are not normally found in a modified/mutated form in normal cells.

15       Within the specific context of prostate cancer, several potentially suitable TAAs have been identified. The first widely studied antigen demonstrated to be over-represented in the prostate gland in cancer modalities was prostatic acid phosphatase (PAP). Arguably, elevated levels of PAP in the bloodstream are considered to be indicative of prostate cancer  
20 (Yam (1974) *Amer J Med* 56:604). Improved methods of cancer detection using this enzyme have been described (WO 79/00475). The biochemical/biophysical structure of the enzyme has also been well studied (Sharief, F.S., et al., (1992) *Biochem Biophys Res Commun* 184:1468-1476; Van Etten, R.L., et al., (1991) *J Biol Chem* 266:9993-9999). Additionally, the  
25 nucleotide sequence encoding human PAP has been determined (Sharief, F.S., et al., (1989) *Biochem Biophys Res Commun* 180:79-86; Tailor, P.G., et al., (1990) *Nucleic Acid Res* 18:4928).

In addition to PAP, other antigen candidates over-represented in prostatic cancer tissue have been identified and/or characterized. The most  
30 prominent among these is human prostate specific antigen (PSA). PSA is a member of the glandular kallikrein family and is a protease with a restricted chymotrypsin-like specificity and is present in the epithelial cells comprising

the prostatic ductal elements. It has been demonstrated in normal prostate and in all primary and metastatic tumor tissue tested, but not in nonprostatic cancer tissues or in normal tissues other than prostate.

The complete amino acid sequence of PSA from human seminal plasma has been determined (Watt K.W. et al. (1986) *Proc Natl Acad Sci USA* 83:3166-3170). PSA consists of a single polypeptide chain with 240 amino acid residues. There is evidence to suggest that the protein is glycosylated. A cDNA encoding PSA has also been isolated (Lundwall, A. and Lilja, H. (1987) *FEBS Lett* 214:317-322; Schultz, P. et al. (1988) *Nucleic Acid Res* 16:6226; Henttu, P. and Bihko, P. (1989) *Biochem and Biophys Res Commun* 160:903-910). The cellular gene for the PSA has also been characterized (Lundwall, A. (1989) *Biochem and Biophys Res Commun* 162:1151-1159, Riegman, P.H.J., et al. (1989) *Biochem and Biophys Res Commun* 159:103-111; Klobeck, G. et al. (1989) *Nucleic Acid Res* 17:3981).

A third prostate-related TAA of potential significance is prostate specific membrane antigen (PSMA). PSMA is similarly found in both benign and neoplastic prostate cells, albeit demonstrating a much greater presence in malignant cells by comparison to benign cells. The presence of PSMA has also been demonstrated on metastatic prostate cells. However, unlike PSA, PSMA is an integral membrane protein. cDNA encoding PSMA has been isolated and characterized (Israeli, R.S. et al. (1993) *Cancer Res* 53:227-230). The cDNA is 2.65 kilobases in length and portions thereof possess significant homology to the nucleic acid sequence coding for human transferrin receptor.

More recently, a number of reports have been published examining the potential usefulness of the prostate-related TAAs in the development of effective cancer immunotherapies. As is well known and documented, T-cell receptors on CD8<sup>+</sup> T cells recognize a complex consisting of peptide (derived from antigen),  $\beta$ -2 microglobulin and class I major histocompatibility complex (MHC) heavy chain (i.e. HLA-A, B, or C in humans). As such, the identification of appropriate MHC class 1-restricted peptides (potentially derived from TAAs) might play an important role in the assessment and/or optimization of


immunotherapeutic treatment strategies. Two important factors must be considered in the assessment of the potential of any TAA epitopes. These are: (1) the peptide must demonstrate an ability to lodge into the peptide binding groove of the respective class 1 MHC molecule and, (2) the peptide must possess sufficiently high binding affinity for the class 1 MHC molecules such that the peptide/class 1 MHC/B2-microglobulin (B2-M)-trimolecular complex displayed on antigen presenting cells is sufficiently stable for presentation, thus resulting in the subsequent activation of the appropriate subset of CD8<sup>+</sup> effector cell.

In view of the foregoing, there is a need in the art to identify and characterize immunogenic peptides derived from PSA (and nucleic acids coding therefor) that can be used to develop effective immunotherapies.

#### **SUMMARY OF THE INVENTION**

The inventors have identified a number of immunogenic peptides derived from prostate-specific antigen (PSA). The PSA derived peptides are useful in treating prostate cancer.

Accordingly, in one embodiment the present invention provides a PSA derived peptide comprising a sequence of the Formula I:



wherein

$n = 0$  or  $1$ ;

each  $X_1$  is independently selected from leucine or methionine;

each  $X_2$  is independently selected from valine or leucine; and

each  $X$  is independently selected from any amino acid,

and fragments, elongations, analogs or derivatives of the PSA derived peptide.

In a preferred embodiment of the invention, the peptides are selected from the group consisting of MWVPVVFL (SEQ ID NO: 1), VLVHPQWVL (SEQ ID NO: 2), and KLQCVDLHV (SEQ ID NO: 3), or a fragment, analog, derivative or elongation of the PSA derived peptide.

In a further aspect, the present invention encompasses nucleic acids coding for the PSA derived peptides. In preferred embodiments, the nucleic

acids have a sequence selected from the group consisting of the sequence of SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9. In yet further embodiments, the invention includes recombinant nucleic acids into which has been inserted a nucleic acid coding for a PSA peptide of the invention.

5       The invention also encompasses compositions. Accordingly, the invention includes compositions of PSA peptides, nucleic acids coding therefore, and/or recombinant nucleic acids into which has been inserted a nucleic acid coding for a PSA peptide of the invention. Said compositions may also comprise suitable adjuvants.

10       Methods of eliciting an immune response in a human are also encompassed by aspects of the invention. Accordingly, methods of eliciting an immune response in an animal comprising the administration of an effective amount of a PSA peptide, a nucleic acid coding therefor, and/or a recombinant nucleic acid into which has been inserted a nucleic acid coding  
15 for PSA peptide, and compositions thereof (with/without an adjuvant) are included within the scope of the invention.

Other features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while  
20 indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from said detailed description.

#### **BRIEF DESCRIPTION OF THE FIGURES**

25       Figure 1 is a graphical representation of effector immune responses elicited in transgenic mice by a PSA peptide.

#### **DETAILED DESCRIPTION OF THE INVENTION**

The following standard one letter and three letter abbreviations for the amino acid residues may be used throughout the specification: A, Ala -  
30 alanine; R, Arg - Arginine; N, Asn - Asparagine; D, Asp - Aspartic acid; C, Cys - Cysteine; Q, Gln - Glutamine; E, Glu - Glutamic acid; G, Gly - Glycine; H, His - Histidine; I, Ile - Isoleucine; L, Leu - Leucine; K, Lys - Lysine; M, Met -


Methionine; F, Phe - Phenylalanine; P, Pro - Proline; S, Ser - Serine; T, Thr - Threonine; W, Trp - Tryptophan; Y, Tyr - Tyrosine; and V, Val - Valine.

### **I. PSA Derived Peptides**

As hereinbefore mentioned, the present inventors have identified and  
5 characterized novel peptides, and nucleic acids encoding these novel peptides, which are derived from prostate-specific antigen (PSA).

Accordingly, the invention provides isolated PSA derived peptides and/or nucleic acids coding therefor which are capable of eliciting an immune response in an animal.

10 In one embodiment, the present invention provides a PSA derived peptide that is capable of eliciting an immune response comprising a sequence of the Formula I:



wherein

15  $n = 0$  or  $1$ ;

each  $X_1$  is independently selected from leucine or methionine;

each  $X_2$  is independently selected from valine or leucine; and

each  $X$  is independently selected from any amino acid;

and fragments, elongations, analogs or derivatives of the PSA derived  
20 peptide.

The term "amino acid" includes all of the naturally occurring amino acids as well as modified amino acids.

Preferred peptides of Formula I include MWVPVFL (SEQ ID NO: 1),  
VLVHPQWWL (SEQ ID NO: 2), and KLQCVDLHV (SEQ ID NO: 3).

25 The phrases "PSA derived peptide(s)" and "PSA peptide(s)" as used herein mean a peptide of Formula I as described above and includes all analogs, derivatives, fragments and elongations thereof which maintain the ability to elicit an immune response in an animal. Preferably, the PSA derived peptide consists essentially of the sequence of the Formula I, more  
30 preferably, the PSA peptides are as shown in SEQ ID NOS:1-3. Collectively, the PSA derived peptides defined herein are referred to as the PSA peptides of the invention.

The term "analog" includes any peptide having an amino acid residue sequence substantially identical to the sequence of the PSA derived peptides shown herein in which one or more residues have been conservatively substituted with a functionally similar residue and which displays the ability to

5 mimic a PSA derived peptide. Examples of conservative substitutions include the substitution of one non-polar (hydrophobic) residue such as alanine, isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between threonine and serine, the

10 substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another. The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that such polypeptide displays the requisite

15 activity.

The term "derivative" refers to a peptide having one or more residues chemically derivatized by reaction of a functional side group. Such derivatized molecules include for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene

20 sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to

25 form N-im-benzylhistidine. Also included as derivatives are those peptides which contain one or more naturally occurring amino acid derivatives of the twenty standard amino acids. For examples: 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be

30 substituted for serine; and ornithine may be substituted for lysine. PSA derived peptides of the present invention also include any peptide having one or more additions and/or deletions or residues relative to the sequence of a



polypeptide whose sequence is shown herein, so long as the requisite activity is maintained or increased.

The term "fragment" refers to any subject peptide having an amino acid residue sequence shorter than that of a PSA peptide of the invention.

- 5 Preferably, the fragment does not contain less than 5, more preferably not less than 8, amino acids.

The term "elongation" refers to any subject peptide having additional amino acid residues added to either end of the peptide, preferably from 1 to 10 amino acid residues, added to either the amino-terminal and/or carboxy-  
10 terminal end of a PSA peptide of the invention.

The invention includes cyclic derivatives of the PSA derived peptides of the invention. Cyclization allows the peptide to assume a more favourable conformation. Cyclization of the peptides may be achieved using techniques known in the art. In particular, disulphide bonds may be formed between two  
15 appropriately spaced components having free sulphhydryl groups. The bonds may be formed between side chains of amino acids, non-amino acid components or a combination of the two.

In a further aspect, the invention includes lipopeptide derivatives of the PSA derived peptides of the invention. Lipopeptides enhance the induction of  
20 CTL responses against antigens in vivo (See e.g. Deres et al., *Nature* 342, 561-564 (1989); Loing et al., *J. Immunol.* 164(2), 900-907 (2000)) and constitute potent adjuvants in parenteral and mucosal immunization (Baier et al., *Immunobiology* 201, 391-405 (2000)). The lipopeptides of the present invention comprise a PSA derived peptide and one or more chains derived  
25 from fatty acids and/or steroid groups, and also include synthetic lipopeptides. The lipopeptides may be prepared using techniques known in the art. In particular, the fatty acids and/or steroid groups may be coupled on the alpha-NH<sub>2</sub> or epsilon-NH<sub>2</sub> functional groups of the amino acid residues of the PSA derived peptide.

30 Peptides of the present invention may be converted into pharmaceutical salts by reacting with inorganic acids including hydrochloric acid, sulphuric acid, hydrobromic acid, phosphoric acid, etc., or organic acids

including formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, succinic acid, malic acid, tartaric acid, citric acid, benzoic acid, salicylic acid, benzenesulphonic acid, and toluenesulphonic acids.

5       The peptides of the invention may be prepared as N-terminal or C-terminal fusion proteins. The fusion proteins may be prepared by fusing, through recombinant techniques or by chemical crosslinking, the N terminal or C-terminal of the peptide, and the sequence of a selected protein or selectable marker with a desired biological function. The resultant fusion  
10 proteins contain the peptide fused to the selected protein or marker protein as described herein. Examples of proteins which may be used to prepare fusion proteins include immunoglobulins, glutathione-S-transferase, hemagglutinin, and truncated myc.

      The peptides of the invention may be used to prepare monoclonal or  
15 polyclonal antibodies. Conventional methods can be used to prepare the antibodies. As to the details relating to the preparation of monoclonal antibodies reference can be made to Goding, J.W., Monoclonal Antibodies: Principles and Practice, 2nd Ed., Academic Press, London, 1986. As discussed below, the antibodies may be used to identify proteins similar or  
20 related to PSA.

      The peptides and antibodies specific for the peptides of the invention may be labelled using conventional methods with various enzymes, fluorescent materials, luminescent materials and radioactive material. Suitable enzymes, fluorescent materials, luminescent materials, and  
25 radioactive material are well known to the skilled artisan. Labelled antibodies specific for the peptides of the invention may be used to screen for proteins similar or related to PSA as discussed in further detail below. Furthermore, the practice of the present invention will employ (unless otherwise indicated) conventional techniques of immunology, molecular biology, cell biology and  
30 recombinant DNA technology which are within the skill of the art.

## **II. Nucleic Acid Molecules of the Invention**

The present invention also includes isolated nucleic acid molecules encoding the PSA derived peptides of the invention.

The term "isolated" refers to a nucleic acid substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors, or other chemicals when chemically synthesized. The term "nucleic acid" is intended to include DNA and RNA and can be either double stranded or single stranded. As such, these nucleic acids comprise the relevant base sequences coding for the aforementioned PSA peptides. For purposes of definitiveness, the "relevant base sequences coding for the aforementioned PSA peptides" further encompasses complementary nucleic acid sequences.

In an embodiment of the invention, isolated nucleic acid molecules are provided having sequences which encode PSA peptides having the amino acid sequences as shown in SEQ ID NOS:1-3.

In a preferred embodiment, the invention provides isolated nucleic acid sequences comprising:

- (a) nucleic acid sequences as shown in SEQ ID NOS:7-9 wherein T can also be U;
- (b) nucleic acid sequences that are complementary to the nucleic acid sequences of (a);
- (c) nucleic acid sequences that have substantial sequence homology to the nucleic acid sequences of (a) or (b);
- (d) nucleic acid sequences that are analogs of the nucleic acid sequences of (a), (b) or (c); or
- (e) nucleic acid sequences that hybridize to the nucleic acid sequences of (a), (b), (c) or (d) under stringent hybridization conditions.

The term "sequence that has substantial sequence homology" means those nucleic acid sequences which have slight or inconsequential sequence variations from the sequences in (a) or (b), i.e., the sequences function in substantially the same manner and can be used to elicit an immune response. The variations may be attributable to local mutations or structural modifications. Nucleic acid sequences having substantial homology include

nucleic acid sequences having at least 65%, more preferably at least 85%, and most preferably 90-95% identity with the nucleic acid sequences as shown in SEQ ID NOS:7-9.

The term "sequence that hybridizes" means a nucleic acid sequence that can hybridize to a sequence of (a), (b), (c) or (d) under stringent hybridization conditions. Appropriate "stringent hybridization conditions" which promote DNA hybridization are known to those skilled in the art, or may be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the following may be employed: 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 x SSC at 50°C; 0.2 x SSC at 50°C to 65°C; or 2.0 x SSC at 44°C to 50°C. The stringency may be selected based on the conditions used in the wash step. For example, the salt concentration in the wash step can be selected from a high stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be at high stringency conditions, at about 65°C.

The term "a nucleic acid sequence which is an analog" means a nucleic acid sequence which has been modified as compared to the sequence of (a), (b) or (c) wherein the modification does not alter the utility of the sequence as described herein. The modified sequence or analog may have improved properties over the sequence shown in (a), (b) or (c). One example of a modification to prepare an analog is to replace one of the naturally occurring bases (i.e. adenine, guanine, cytosine or thymidine) of the sequences shown in SEQ ID NOS:7-9, with a modified base such as such as xanthine, hypoxanthine, 2-aminoadenine, 6-methyl, 2-propyl and other alkyl adenines, 5-halo uracil, 5-halo cytosine, 6-aza uracil, 6-aza cytosine and 6-aza thymine, pseudo uracil, 4-thiouracil, 8-halo adenine, 8-aminoadenine, 8-thiol adenine, 8-thiolalkyl adenines, 8-hydroxyl adenine and other 8-substituted adenines, 8-halo guanines, 8 amino guanine, 8-thiol guanine, 8-thiolalkyl guanines, 8-hydroxyl guanine and other 8-substituted guanines, other aza and deaza uracils, thymidines, cytosines, adenines, or guanines, 5-trifluoromethyl uracil and 5-trifluoro cytosine.

Another example of a modification is to include modified phosphorous or oxygen heteroatoms in the phosphate backbone, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages in the nucleic acid molecules shown in SEQ ID NOS:7-9.

- 5 For example, the nucleic acid sequences may contain phosphorothioates, phosphotriesters, methyl phosphonates, and phosphorodithioates.

A further example of an analog of a nucleic acid molecule of the invention is a peptide nucleic acid (PNA) wherein the deoxyribose (or ribose) phosphate backbone in the DNA (or RNA), is replaced with a polyamide backbone which is similar to that found in peptides (P.E. Nielsen, et al  
10 Science 1991, 254, 1497). PNA analogs have been shown to be resistant to degradation by enzymes and to have extended lives *in vivo* and *in vitro*. PNAs also bind stronger to a complimentary DNA sequence due to the lack of charge repulsion between the PNA strand and the DNA strand. Other nucleic  
15 acid analogs may contain nucleotides containing polymer backbones, cyclic backbones, or acyclic backbones. For example, the nucleotides may have morpholino backbone structures (U.S. Pat. No. 5,034,506). The analogs may also contain groups such as reporter groups, a group for improving the pharmacokinetic or pharmacodynamic properties of nucleic acid sequences.

- 20 It will be appreciated that the invention includes nucleic acid molecules encoding elongations of peptides of the invention, and analogs and homologs of peptides of the invention and elongations thereof, as described above.

Isolated and purified nucleic acid molecules having sequences which differ from the nucleic acid sequence of the invention due to degeneracy in  
25 the genetic code are also within the scope of the invention. Such nucleic acids encode functionally equivalent peptides but differ in sequence from the above mentioned sequences due to degeneracy in the genetic code.

An isolated nucleic acid molecule of the invention which comprises DNA can be isolated by preparing a labelled nucleic acid probe based on all  
30 or part of the nucleic acid sequences of the invention and using this labelled nucleic acid probe to screen an appropriate DNA library (e.g. a cDNA or genomic DNA library). For example, a genomic library isolated can be used

to isolate a DNA encoding a novel protein of the invention by screening the library with the labelled probe using standard techniques. Nucleic acids isolated by screening of a cDNA or genomic DNA library can be sequenced by standard techniques.

- 5 An isolated nucleic acid molecule of the invention which is DNA can also be isolated by selectively amplifying a nucleic acid encoding a novel peptide of the invention using the polymerase chain reaction (PCR) methods and cDNA or genomic DNA. It is possible to design synthetic oligonucleotide primers from the nucleic acid sequence of the invention for use in PCR. A
- 10 nucleic acid can be amplified from cDNA or genomic DNA using these oligonucleotide primers and standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. It will be appreciated that cDNA may be prepared from mRNA, by isolating total cellular mRNA by a variety of
- 15 techniques, for example, by using the guanidinium-thiocyanate extraction procedure of Chirgwin et al., Biochemistry, 18, 5294-5299 (1979). cDNA is then synthesized from the mRNA using reverse transcriptase (for example, Moloney MLV reverse transcriptase available from Gibco/BRL, Bethesda, MD, or AMV reverse transcriptase available from Seikagaku America, Inc., St.
- 20 Petersburg, FL).

- An isolated nucleic acid molecule of the invention which is RNA can be isolated by cloning a cDNA encoding a novel peptide of the invention into an appropriate vector which allows for transcription of the cDNA to produce an RNA molecule which encodes a peptide of the invention. For example, a
- 25 cDNA can be cloned downstream of a bacteriophage promoter, (e.g., a T7 promoter) in a vector, cDNA can be transcribed *in vitro* with T7 polymerase, and the resultant RNA can be isolated by standard techniques.

- A nucleic acid molecule of the invention may also be chemically synthesized using standard techniques. Various methods of chemically
- 30 synthesizing polydeoxynucleotides are known, including solid-phase synthesis which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See e.g., Itakura et al. U.S. Patent

No. 4,598,049; Caruthers et al. U.S. Patent No. 4,458,066; and Itakura U.S. Patent Nos. 4,401,796 and 4,373,071).

Determination of whether a particular nucleic acid molecule encodes a novel peptide of the invention may be accomplished by expressing the cDNA  
5 in an appropriate host cell by standard techniques, and testing the activity of the protein using the methods as described herein. A cDNA having the activity of a novel peptide of the invention so isolated can be sequenced by standard techniques, such as dideoxynucleotide chain termination or Maxam-Gilbert chemical sequencing, to determine the nucleic acid sequence and the  
10 predicted amino acid sequence of the encoded peptide.

The sequence of a nucleic acid molecule of the invention may be inverted relative to its normal presentation for transcription to produce an antisense nucleic acid molecule which are more fully described herein. Preferably, an antisense sequence is constructed by inverting a region  
15 preceding the initiation codon or an unconserved region. In particular, the nucleic acid sequences contained in the nucleic acid molecules of the invention or a fragment thereof, may be inverted relative to its normal presentation for transcription to produce antisense nucleic acid molecules.

The invention also provides nucleic acids encoding fusion proteins  
20 comprising a novel peptide of the invention and a selected protein, or a selectable marker protein (see below).

### **III. Preparation of the PSA Derived Peptides**

The PSA peptides of the invention may be prepared using a variety of methods known to one skilled in the art. Accordingly, PSA peptides may be  
25 prepared by chemical synthesis using techniques well known in the chemistry of proteins such as solid phase synthesis (Merrifield, J. Am. Chem. Assoc. 85:2149-2154 (1964)) or synthesis in homogenous solution (Houbenweyl, Methods of Organic Chemistry, ed. E. Wansch, Vol. 15, pts. I and II, Thieme, Stuttgart (1987)).

30 The peptides of the invention may also be produced by recombinant DNA technology. To prepare the peptides of the invention by recombinant DNA techniques, a DNA sequence encoding the PSA derived peptide must

be prepared. Consequently, the present invention also provides purified and isolated nucleic acids having a nucleotide sequence coding for PSA derived peptides. In one embodiment of the invention, the nucleic acid has a sequence encoding a PSA derived peptide comprising an amino acid  
5 sequence of the formula I wherein the substituents are described above. In other embodiments, the DNA sequence encoding a PSA derived peptide of the formula I comprises a nucleotide sequence of ATG TGG GTC CCG GTT GTC TTC CTC (SEQ ID NO:7), GTT CTG GTG CAC CCC CAG TGG GTC CTC (SEQ ID NO:8), or AAA CTT CAG TGT GTG GAC CTC CAT GTT (SEQ  
10 ID NO:9).

The present invention also provides an expression vector comprising a DNA molecule encoding a PSA derived peptide adapted for transfection or transformation of a host cell. The nucleic acid molecules of the present invention may be incorporated in a known manner into an appropriate  
15 expression vector which ensures expression of the protein. Possible expression vectors include but are not limited to cosmids, plasmids, or modified viruses (e.g. replication defective retroviruses, adenoviruses and adeno-associated viruses). The vector should be compatible with the host cell used. The expression vectors are "suitable for transformation of a host cell", which means that the expression vectors contain a nucleic acid molecule of the invention and regulatory sequences selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid molecule. "Operatively linked" is intended to mean that the nucleic acid is linked to regulatory sequences in a manner which allows  
20 expression of the nucleic acid.

The invention therefore contemplates a recombinant expression vector containing a nucleic acid molecule of the invention, or a fragment thereof, and the necessary regulatory sequences for the transcription and translation of the inserted protein-sequence.

Suitable regulatory sequences may be derived from a variety of  
30 sources, including bacterial, fungal, viral, mammalian, or insect genes (For example, see the regulatory sequences described in Goeddel, Gene



Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA 1990). Selection of appropriate regulatory sequences is dependent on the host cell chosen as discussed below, and may be readily accomplished by one of ordinary skill in the art. Examples of such regulatory

- 5 sequences include: a transcriptional promoter and enhancer or RNA polymerase binding sequence, a ribosomal binding sequence, including a translation initiation signal. Additionally, depending on the host cell chosen and the vector employed, other sequences, such as an origin of replication, additional DNA restriction sites, enhancers, and sequences conferring
- 10 inducibility of transcription may be incorporated into the expression vector. It will also be appreciated that the necessary regulatory sequences may be supplied by the native A and B chains and/or its flanking regions.

- The recombinant expression vectors of the invention may also contain a marker gene which facilitates the selection of host cells transformed or
- 15 transfected with a recombinant molecule of the invention. Examples of selectable marker genes are genes encoding a protein such as G418 and hygromycin which confer resistance to certain drugs,  $\beta$ -galactosidase, chloramphenicol acetyltransferase, firefly luciferase, or an immunoglobulin or portion thereof such as the Fc portion of an immunoglobulin preferably IgG.
- 20 Transcription of the selectable marker gene is monitored by changes in the concentration of the selectable marker protein such as  $\beta$ -galactosidase, chloramphenicol acetyltransferase, or firefly luciferase. If the selectable marker gene encodes a protein conferring antibiotic resistance such as neomycin resistance transformant cells can be selected with G418. Cells that
- 25 have incorporated the selectable marker gene will survive, while the other cells die. This makes it possible to visualize and assay for expression of recombinant expression vectors of the invention and in particular to determine the effect of a mutation on expression and phenotype. It will be appreciated that selectable markers can be introduced on a separate vector from the
- 30 nucleic acid of interest.

Recombinant expression vectors can be introduced into host cells to produce a transformant host cell. The term "transformant host cell" is

intended to include prokaryotic and eukaryotic cells which have been transformed or transfected with a recombinant expression vector of the invention. The terms "transformed with", "transfected with", "transformation" and "transfection" are intended to encompass introduction of nucleic acid (e.g. a vector) into a cell by one of many possible techniques known in the art. Prokaryotic cells can be transformed with nucleic acid by, for example, electroporation or calcium-chloride mediated transformation. Nucleic acid can be introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride co-precipitation, DEAE-dextran mediated transfection, lipofectin, electroporation or microinjection. Suitable methods for transforming and transfecting host cells can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks.

Suitable host cells include a wide variety of prokaryotic and eukaryotic host cells. For example, the proteins of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus), yeast cells or mammalian cells. Other suitable host cells can be found in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1991).

More particularly, bacterial host cells suitable for carrying out the present invention include *E. coli*, *B. subtilis*, *Salmonella typhimurium*, and various species within the genus' *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, as well as many other bacterial species well known to one of ordinary skill in the art. Suitable bacterial expression vectors preferably comprise a promoter which functions in the host cell, one or more selectable phenotypic markers, and a bacterial origin of replication. Representative promoters include the  $\beta$ -lactamase (penicillinase) and lactose promoter system (see Chang et al., *Nature* 275:615 (1978)), the *trp* promoter (Nichols and Yanofsky, *Meth. in Enzymology* 101:155, 1983) and the *tac* promoter (Russell et al., *Gene* 20: 231, 1982). Representative selectable markers include various antibiotic resistance markers such as the kanamycin or ampicillin resistance genes. Suitable expression vectors include but are not

limited to bacteriophages such as lambda derivatives or plasmids such as pBR322 (see Bolivar et al., Gene 2:95, (1977)), the pUC plasmids pUC18, pUC19, pUC118, pUC119 (see Messing, Meth in Enzymology 101:20-77, 1983 and Vieira and Messing, Gene 19:259-268 (1982)), and pNH8A, 5 pNH16a, pNH18a, and Bluescript M13 (Stratagene, La Jolla, Calif.). Typical fusion expression vectors which may be used are discussed above, e.g. pGEX (Amrad Corp., Melbourne, Australia), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ). Examples of inducible non-fusion expression vectors include pTrc (Amann et al., Gene 69:301-315 10 (1988)) and pET 11d (Studier et al., Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, California, 60-89 (1990)).

Yeast and fungi host cells suitable for carrying out the present invention include, but are not limited to *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, the genera *Pichia* or *Kluyveromyces* and 15 various species of the genus *Aspergillus*. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari. et al., Embo J. 6:229-234 (1987)), pMFa (Kurjan and Herskowitz, Cell 30:933-943 (1982)), pJRY88 (Schultz et al., Gene 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA). Protocols for the transformation of yeast and fungi are well 20 known to those of ordinary skill in the art.(see Hinnen et al., Proc. Natl. Acad. Sci. USA 75:1929 (1978); Itoh et al., J. Bacteriology 153:163 (1983), and Cullen et al. (Bio/Technology 5:369 (1987)).

Mammalian cells suitable for carrying out the present invention include, among others: COS (e.g., ATCC No. CRL 1650 or 1651), BHK (e.g. ATCC 25 No. CRL 6281), CHO (ATCC No. CCL 61), HeLa (e.g., ATCC No. CCL 2), 293 (ATCC No. 1573) and NS-1 cells. Suitable expression vectors for directing expression in mammalian cells generally include a promoter (e.g., derived from viral material such as polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40), as well as other transcriptional and translational control 30 sequences. Examples of mammalian expression vectors include pCDM8 (Seed, B., Nature 329:840 (1987)) and pMT2PC (Kaufman et al., EMBO J. 6:187-195 (1987)).

Given the teachings provided herein, promoters, terminators, and methods for introducing expression vectors of an appropriate type into plant, avian, and insect cells may also be readily accomplished. For example, within one embodiment, the proteins of the invention may be expressed from  
5 plant cells (see Sinkar et al., J. Biosci (Bangalore) 11:47-58 (1987), which reviews the use of *Agrobacterium rhizogenes* vectors; see also Zambryski et al., Genetic Engineering, Principles and Methods, Hollaender and Setlow (eds.), Vol. VI, pp. 253-278, Plenum Press, New York (1984), which describes the use of expression vectors for plant cells, including, among others,  
10 pAS2022, pAS2023, and pAS2034).

Insect cells suitable for carrying out the present invention include cells and cell lines from *Bombyx* or *Spodoptera* species. Baculovirus vectors available for expression of proteins in cultured insect cells (SF 9 cells) include the pAc series (Smith et al., Mol. Cell Biol. 3:2156-2165 (1983)) and the pVL  
15 series (Lucklow, V.A., and Summers, M.D., Virology 170:31-39 (1989)). Some baculovirus-insect cell expression systems suitable for expression of the recombinant proteins of the invention are described in PCT/US/02442.

The recombinant expression vectors containing the nucleotide sequences encoding the PSA derived peptides may also contain genes which  
20 encode a fusion moiety (i.e. a "fusion protein") which provides increased expression of the recombinant peptide; increased solubility of the recombinant peptide; and aid in the purification of the target recombinant peptide by acting as a ligand in affinity purification. For example, a proteolytic cleavage site may be added to the target recombinant protein to allow separation of the  
25 recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Typical fusion expression vectors include pGEX (Amrad Corp., Melbourne, Australia), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the recombinant  
30 protein. By way of illustration, the DNA sequence encoding an enhanced affinity PSA derived peptide may be cloned into a pGEX-type plasmid for co-expression with a 26 kD protein glutathione-S-transferase (GST): pGEX-2T,

pGEX-2TK, pGEX-3X, pGEX-1T, pGEX-4T, pGEX-5X. Said plasmids are transformed into Escherichia coli HB101 cells, and positive clones can then be selected by standard hybridization techniques identifying GST-PSA peptide fusion protein. Colonies with the pGEX-2T plasmids containing DNA  
5 encoding an enhanced affinity PSA peptide may be grown in 5 mL of LB/amp medium at 37°C for overnight. Isopropylthio--D-galactoside is added to the culture to a final concentration of 0.1 mM to induce the biosynthesis of the fusion protein. The cultures are grown for 90 min post induction and the cells are harvested by centrifugation and lysed in 1 mL of 50 mM Tris-HCl, pH 8.0,  
10 containing 2 M urea and 1% Triton X-100. Cells can then be further disrupted by sonication and centrifuged at 15,000 x g for 20 min to separate soluble from insoluble fractions. The supernatant containing the GST-PSA peptide fusion protein is then used for extraction of the fusion protein by standard procedures (GST Purification Module: Pharmacia Cat. #27-4570-01, -02).  
15 Subsequently, the fusion protein may be treated with an enzyme to release the enhanced affinity PSA peptide (thrombin for pGEX-2T, pGEX-2TK, pGEX-1T, pGEX-4T; factor Xa for pGEX-3X and pGEX-5X).

Additional embodiments of the invention encompass recombinant nucleic acids further comprising inserts. These inserts code for the PSA  
20 peptides hereinbefore described. Further embodiments encompass recombinant nucleic acids wherein the insert comprises a sequence chosen from the group consisting of SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9. As defined herein, "recombinant nucleic acids" encompass (but is not limited to) viruses, bacterial DNA, naked/free DNA and RNA.

25 **IV. Applications of the Peptides and Nucleic Acids**

The inventors have demonstrated that the PSA derived peptides of the present invention are immunogenic and capable of eliciting immune responses *in vivo*. Consequently, the present invention includes the use of one or more PSA derived peptides of the invention to modulate immune  
30 responses. Accordingly, the present invention provides a method of modulating immune responses comprising administering an effective amount

of a PSA derived peptide or a nucleic acid molecule encoding a PSA peptide of the invention to a cell or animal in need thereof.

The term "animal" as used herein includes all members of the animal kingdom including mammals, preferably humans.

5       The term "eliciting an immune response" is defined as initiating, triggering, causing, enhancing, improving or augmenting any response of the immune system, for example, of either a humoral or cell-mediated nature. The initiation or enhancement of an immune response can be assessed using assays known to those skilled in the art including, but not limited to, antibody  
10       assays (for example ELISA assays), antigen specific cytotoxicity assays and the production of cytokines (for example ELISPOT assays). Preferably, the peptides and nucleic acids of the present invention, and the method of the present invention trigger or enhance a cellular immune response, more preferably a cytotoxic T cell response.

15       The term "effective amount" as used herein means an amount effective, at dosages and for periods of time necessary to achieve desired results.

More specifically, the peptides of the invention may be used in the prophylaxis or treatment of pathological conditions such as cancer, including  
20       tumor metastasis, in a mammal.

Additional embodiments of the invention encompass compositions comprising PSA peptides, and/or nucleic acids coding for PSA peptides, and /or recombinant nucleic acids into which has been inserted a nucleic acid sequence coding for PSA peptide(s) (all of which have been herein before  
25       described). The peptides and nucleic acid molecules may be formulated into pharmaceutical compositions for administration to subjects in a biologically compatible form suitable for administration. By biologically compatible form suitable for administration is meant a form of the substance to be administered in which any toxic effects are outweighed by the therapeutic  
30       effects. The substances may be administered to living organisms including humans, and animals in a therapeutically effective amount. Administration of an effective amount of the pharmaceutical compositions of the present

invention is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, a therapeutically active amount of a substance may vary according to factors such as the disease state, age, sex and weight of the individual, and the ability of peptide to elicit a desired response in the individual. Dosage regime may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

The active substance may be administered in a convenient manner such as by topical or transdermal application, injection (subcutaneous, intravenous, etc.), oral administration, inhalation, or rectal administration. Depending on the route of administration, the active substance may be coated in a material to protect the compound from the action of enzymes, acids and other natural conditions which may inactivate the compound.

Several modes of administration are available when using a composition containing a nucleic acid molecule encoding a PSA derived peptide of the invention. Recombinant molecules comprising a nucleic acid sequence encoding a PSA derived protein (as described above), or fragment thereof, may be directly introduced into cells or tissues *in vivo* using delivery vehicles such as retroviral vectors, adenoviral vectors and DNA virus vectors. They may also be introduced into cells *in vivo* using physical techniques such as microinjection and electroporation or chemical methods such as coprecipitation and incorporation of DNA into liposomes. Recombinant molecules may also be delivered in the form of an aerosol or by lavage. The nucleic acid molecules of the invention may also be applied extracellularly such as by direct injection into cells.

The compositions described herein can be prepared by *per se* known methods for the preparation of pharmaceutically acceptable compositions which can be administered to subjects, such that an effective quantity of the active substance is combined in a mixture with a pharmaceutically acceptable vehicle. Suitable vehicles are described, for example, in Remington's Pharmaceutical Sciences (Remington's Pharmaceutical Sciences (1985),

Mack Publishing Company, Easton, Pa., USA). On this basis, the compositions include, albeit not exclusively, solutions of the substances in association with one or more pharmaceutically acceptable vehicles or diluents, and may be contained in buffered solutions with a suitable pH and/or  
5 be iso-osmotic with physiological fluids. In this regard, reference can be made to U.S. Patent No. 5,843,456.

Compositions for injection include, albeit not exclusively, the peptides or nucleic acids in association with one or more pharmaceutically acceptable vehicles or diluents, and contained in buffered solutions with a suitable pH  
10 and iso-osmotic with the physiological fluids. Any pharmaceutically suitable diluent can be used in the composition for injections: distilled water, physiological or a salt solution, and/or a buffer solution. The composition for injections may be prepared by conventional volume-weight procedures. A certain amount of the peptide is diluted to the necessary volume with a diluent  
15 or solvent. The solution is then filtered through sterilized filters, bottled or ampouled. The resultant solution is a stable transparent liquid, and does not contain any chemical or other impurities.

Solid form preparations for oral administration can be made in the form of tablets, powders, or capsules. It may contain a medium for the active  
20 substance and other additives, including dyes, aromas, etc.

The compositions and treatments are indicated as therapeutic agents or treatments either alone or in conjunction with other therapeutic agents or other forms of treatment.

Immunogenicity can be significantly improved if the immunizing  
25 agent(s) (i.e. PSA peptide, and/or nucleic acid coding therefor, recombinant nucleic acids) and/or composition is, regardless of administration format, co-immunized with an adjuvant. Commonly, adjuvants are used as an 0.05 to 1.0 percent solution in phosphate - buffered saline. Adjuvants enhance the immunogenicity of an immunogen but are not necessarily immunogenic  
30 themselves. Adjuvants may act by retaining the immunogen locally near the site of administration to produce a depot effect facilitating a slow, sustained release of immunogen to cells of the immune system. Adjuvants can also



attract cells of the immune system to an immunogen depot and stimulate such cells to elicit immune responses. As such, embodiments of this invention encompass compositions further comprising adjuvants.

Adjuvants have been used for many years to improve the host immune responses to, for example, vaccines. Intrinsic adjuvants (such as lipopolysaccharides) normally are the components of killed or attenuated bacteria used as vaccines. Extrinsic adjuvants are immunomodulators which are typically non-covalently linked to antigens and are formulated to enhance the host immune responses. Thus, adjuvants have been identified that enhance the immune response to antigens delivered parenterally. Some of these adjuvants are toxic, however, and can cause undesirable side-effects making them unsuitable for use in humans and many animals. Indeed, only aluminum hydroxide and aluminum phosphate (collectively commonly referred to as alum) are routinely used as adjuvants in human and veterinary vaccines. The efficacy of alum in increasing antibody responses to diphtheria and tetanus toxoids is well established. Notwithstanding, it does have limitations. For example, alum is ineffective for influenza vaccination and inconsistently elicits a cell mediated immune response with other immunogens. The antibodies elicited by alum-adjuvanted antigens are mainly of the IgG1 isotype in the mouse, which may not be optimal for protection by some vaccinal agents.

A wide range of extrinsic adjuvants can provoke potent immune responses to immunogens. These include saponins complexed to membrane protein antigens (immune stimulating complexes), pluronic polymers with mineral oil, killed mycobacteria and mineral oil, Freund's complete adjuvant, bacterial products such as muramyl dipeptide (MDP) and lipopolysaccharide (LPS), as well as lipid A, and liposomes.

In one aspect of this invention, adjuvants useful in any of the embodiments of the invention described herein are as follows. Adjuvants for parenteral immunization include aluminum compounds (such as aluminum hydroxide, aluminum phosphate, and aluminum hydroxy phosphate). The antigen can be precipitated with, or adsorbed onto, the aluminum compound

according to standard protocols. Other adjuvants such as RIBI (ImmunoChem, Hamilton, MT) can also be used in parenteral administration.

Adjuvants for mucosal immunization include bacterial toxins (e.g., the cholera toxin (CT), the *E. coli* heat-labile toxin (LT), the *Clostridium difficile* toxin A and the pertussis toxin (PT), or combinations, subunits, toxoids, or mutants thereof). For example, a purified preparation of native cholera toxin subunit B (CTB) can be of use. Fragments, homologs, derivatives, and fusion to any of these toxins are also suitable, provided that they retain adjuvant activity. Preferably, a mutant having reduced toxicity is used. Suitable mutants have been described (e.g., in WO 95/17211 (Arg-7-Lys CT mutant), WO 96/6627 (Arg-192-Gly LT mutant), and WO 95/34323 (Arg-9-Lys and Glu-129-Gly PT mutant)). Additional LT mutants that can be used in the methods and compositions of the invention include, for example Ser-63-Lys, Ala-69-Gly, Glu-110-Asp, and Glu-112-Asp mutants. Other adjuvants (such as a bacterial monophosphoryl lipid A (MPLA) of various sources (e.g., *E. coli*, *Salmonella minnesota*, *Salmonella typhimurium*, or *Shigella flexneri*, saponins, or polylactide glycolide (PLGA) microspheres) can also be used in mucosal administration.

Adjuvants useful for both mucosal and parenteral immunization include polyphosphazene (for example, WO 95/2415), DC-chol (3 b-(N-(N',N'-dimethyl aminomethane)-carbamoyl) cholesterol (for example, U.S. Patent No. 5,283,185 and WO 96/14831) and QS-21 (for example, WO 88/9336).

An animal may be immunized with a PSA peptide(s), a nucleic acid coding therefor, a recombinant nucleic acid wherein a nucleic acid coding for a PSA peptide is inserted therein, and/or a composition of this invention by any conventional route as is known to one skilled in the art. This may include, for example, immunization via a mucosal (e.g., ocular, intranasal, oral, gastric, pulmonary, intestinal, rectal, vaginal, or urinary tract) surface, via the parenteral (e.g., subcutaneous, intradermal, intramuscular, intravenous, or intraperitoneal) route or intranodally. Preferred routes depend upon the choice of the immunogen as will be apparent to one skilled in the art. The administration can be achieved in a single dose or repeated at intervals. The

appropriate dosage depends on various parameters understood by skilled artisans such as the immunogen itself (i.e. peptide vs. nucleic acid (and more specifically type thereof)), the route of administration and the condition of the animal to be vaccinated (weight, age and the like).

- 5           Accordingly, embodiments of this invention encompass methods of eliciting immune responses in animals comprising administering effective amounts of PSA peptide(s), and/or nucleic acid(s) coding therefore, and/or recombinant nucleic acid(s) wherein a nucleic acid coding for a PSA peptide is inserted therein, and/or compositions of the invention. The present
- 10 invention also includes methods of treating cancer comprising administering effective amounts of PSA peptide(s), and/or nucleic acid(s) coding therefore, and/or recombinant nucleic acid(s) wherein a nucleic acid coding for a PSA peptide is inserted therein, and/or compositions of the invention. In a preferred embodiment, the methods of the invention are utilized to treat
- 15 prostate cancer.

- A further embodiment of this invention encompasses a use of effective amounts of PSA peptide(s), and/or nucleic acid(s) coding therefore, and/or recombinant nucleic acid(s) wherein a nucleic acid coding for a PSA peptide is inserted therein, and/or a composition of the present invention to elicit an
- 20 immune response in an animal preferably to treat cancer, more preferably prostate cancer. The present invention further includes a use of effective amounts of PSA peptide(s), and/or nucleic acid(s) coding therefore, and/or recombinant nucleic acid(s) wherein a nucleic acid coding for a PSA peptide is inserted therein, and/or a composition of the present invention to prepare a
- 25 medicament to elicit an immune response in animal, preferably to treat cancer, more preferably prostate cancer.

The following non-limiting examples are illustrative of the present invention:

#### **EXAMPLES**

30   **Experimental**

The analysis of peptides isolated from human class 1 MHC complexes have revealed that peptides which are favored to bind and lodge into the

peptide-binding groove of the human MHC class 1 molecule (such as HLA-A0201) are typically 9 amino acids long (however, peptides of 8-13 amino acids have also been observed). In the majority of cases, these nonamers contain two anchor residues; the first proximal to the amino-(N-) terminus, and the second associated with the carboxy-(C-) terminus. These anchor residues interact with the respective 'pockets' of the peptide-binding groove of the MHC molecules. The amino-(N-) associated anchor residue is typically leucine (L) or methionine (M); the carboxy-(C-) associated residue is typically leucine (L) or valine (V).

Methods of peptide synthesis, cell culture propagation and assays of cytotoxicity are amply reported in the scientific literature and are well within the scope of those skilled in the art; as such, detailed particulars of these methods will not be explicitly discussed.

#### **Example 1**

A number of PSA derived peptide sequences were identified that correspond to the above-mentioned Formula I. These are outlined in Table 1.

#### **Peptide Synthesis**

Solid phase peptide syntheses were conducted on an ABI 430A automated peptide synthesizer according to the manufacturer's standard protocols. The peptides were cleaved from the solid support by treatment with liquid hydrogen fluoride in the presence of thiocresole, anisole, and methyl sulfide. The crude products were extracted with trifluoroacetic acid (TFA) and precipitated with diethyl ether. All peptides were stored in lyophilized form at -20°C.

The peptides synthesized were:

CLP313	MWWPVVFL	(SEQ ID NO: 1)
CLP314	VLVHPQWWL	(SEQ ID NO: 2)
CLP316	KLQCVDLHV	(SEQ ID NO: 3)
CLP315	DLPTQEPAL	(SEQ ID NO: 4)
CLP317	DLHVISNDV	(SEQ ID NO: 5)
CLP320	PLYDMSLLK	(SEQ ID NO: 6)

Prior to immunization of animals, peptides were dissolved in 100% Dimethylsulphoxide (DMSO), subsequently diluted with sterile distilled water (DW) to a final DMSO concentration 5.0%, and stored at  $-20^{\circ}\text{C}$  pending use.

### Example 2

#### 5 Nucleic Acid Sequences Coding for PSA Derived Peptides

The nucleic acid sequence coding for the identified PSA peptides (i.e. SEQ ID. NOs:1-6) were deduced by reference to the nucleic acid sequence disclosed in the publications of Henttu and Vihko ((1989) *Biochem and Biophys Res Commun* 160:903-910), Lundwall and Lilja ((1987) *FEBS Lett.* 214:317-322) and Schulz et al. ((1988) *Nucleic Acids Research* 16:6226).

The coding strand nucleic acid sequences were:

<u>Peptide</u>	<u>Nucleic Acid Sequence</u>	
CLP313(SEQ ID NO:1)	ATGTGGGTCCCGTTGTCTCTCTC	(SEQ ID NO:7)
CLP314(SEQ ID NO:2)	GTTCTGGTGCACCCCCAGTGGGTCTC	(SEQ ID NO:8)
CLP316(SEQ ID NO:3)	AAACTTCAGTGTGTGGACCTCCATGTT	(SEQ ID NO:9)
CLP315(SEQ ID NO:4)	GACCTGCCCACCCAGGAGCCAGCACTG	(SEQ ID NO:10)
CLP317(SEQ ID NO:5)	GACCTCCATGTTATTTCCAATGACGTG	(SEQ ID NO:11)
CLP320(SEQ ID NO:6)	CCGCTCTACGATATGAGCTCTCTGAAG	(SEQ ID NO:12)

### Example 3

#### 15 HLA-A0201 Binding of PSA Derived Peptides

The ability of the PSA derived peptides described in Table 1 to stabilize membrane-bound HLA-A0201 molecule was assessed utilizing the T2 cell line (Dr. Peter Creswell, Yale University). The cell line has been well documented to have a defective TAP (i.e. Transporter for Antigen Processing) transporter function. As a result, the majority of intracellularly generated peptides are not transported into the endoplasmic reticulum and thus are unable to associate with newly synthesized HLA class 1 MHC molecules (i.e. HLA-A0201; Salter, R D and Creswell, P. (1986) *EMBO J* 5:943). The majority of the HLA-A0201

molecules displayed on the surface of T2 cells are therefore empty (contain no peptides) and unstable. The stability of the surface HLA-A0201 molecules can be restored upon interaction with suitable exogenous peptides. The stabilization of the conformation of the class 1 MHC molecules is accompanied by the formation of an immunodominant epitope recognized by a mouse monoclonal antibody (designated BB7.2; American Type Culture Collection (ATCC)). Thus, the detection of this specific epitope is indicative of stable membrane-bound HLA-A0201 molecules loaded with peptide. Subsequent dissociation of peptides from the HLA class 1 MHC molecules results in the loss of BB7.2 monoclonal antibody binding.

T2 cells were propagated in Iscove's complete medium (Iscove's medium supplemented with 10% heat-inactivated bovine serum, 120.0 units per ml of penicillin G sodium, 120 µg per ml of streptomycin sulphate, and 0.35 mg per ml of L-glutamine). The ability of PSA peptides (Table 1) to bind and stabilize surface HLA-A0201 molecules on T2 cells was determined utilizing a protocol documented in the art (Deng, Y. (1997) *J Immunol* 158:1507-1515). In essence,  $0.5 \times 10^6$  T2 cells were incubated with 125.0 µM of the test peptide in 200.0 µl of serum-free culture medium (Iscove's medium supplemented with 120.0 units per ml of penicillin G sodium, 120.0 µg per ml of streptomycin sulphate and 0.35 mg per ml of L-glutamine) overnight at 37°C. Subsequently, cells were washed twice with Iscove's medium (without bovine serum) to remove free PSA peptide. To the cell pellet was added 1.0 ml of complete Iscove's medium (supplemented with fetal bovine serum (10% final)) containing 5.0 µg per ml of brefeldin A (Sigma), 12.5 µg per ml of anisomycin (Sigma) and 5.0 µg per ml of cyclohexamide (Sigma). The samples were incubated for 3.0 hr in a 37°C CO<sub>2</sub> incubator. The cells were subsequently incubated 30 min on ice prior to two washes with ice-cold PBA (phosphate buffered saline (pH 7.2) containing 2.0% bovine serum). 100.0 µl of PBA containing 5.0 µg of monoclonal antibody BB7.2 was added to each test sample. The reaction was allowed to proceed on ice for 45 min. Cells were then washed three times with ice-cold PBA. The binding of BB7.2 was detected via the addition of 100.0 µl of PBA containing 1.0 µg of goat anti-

mouse IgG-Fc fluorescein (FITC) conjugate (BETHYL Laboratories Inc.). After a 30 min incubation on ice, cells were twice washed with ice-cold PBA, and twice with ice-cold PBS (pH 7.2). Cells were immediately fixed by the addition of 50.0  $\mu$ l of 1.0% paraformaldehyde. Samples were analyzed by Flow Cytometry. Results were expressed in units of Fluorescence Index (FI), calculated by the equation:

$$\frac{\text{Mean Fluorescence (MF) of experimental sample (peptide treated)} - \text{MF of control sample (cells not peptide treated)}}{\text{MF of control sample (cells not peptide treated)}}$$

An FI value of 0.5 or greater was deemed to be significant.

The results depicted in Table 1 reveal that a negative control peptide (CLP-320) containing a binding motif for the gene product of HLA-A1 does not exhibit binding to the membrane-bound HLA-A0201 molecules on T2 cells. Of the 6 PSA peptides tested, 3 (i.e. CLP-313, CPL-314, CPL-316) demonstrated HLA-A0201 binding as judged by Fluorescence Index (FI) values (Table 1).

#### **Example 4**

#### **Immunogenicity of PSA Derived Peptides**

The A2Kb transgenic mouse was used to assess the immunogenicity of the HLA-A0201 binding PSA peptides. Mice of the B10 background (transgenic for the A2Kb chimeric gene) were purchased from the Scripps Clinic in California, USA. Mice were injected subcutaneously at the base of the tail with a dose of inoculum prepared by emulsifying 100.0  $\mu$ g of the test peptide and 100.0  $\mu$ g of an I-A<sup>b</sup>-restricted peptide (described in Milich, D.R. et al. (1987) *J. Immunol* 139: 1223-1231) in incomplete Freund's adjuvant (IFA). Spleens of the experimental animals were collected on the 10th or 11th day post immunization. Spleenocytes of the experimental mice were prepared and cultured to enrich for CTLs before being assessed for effector activity. *In vitro* re-stimulation of the *in vivo* generated CTLs was performed by co-culturing in a 25 cm<sup>2</sup> tissue culture flask 3 x 10<sup>7</sup> responder cells (i.e. spleenocytes) with

1.3 x 10<sup>7</sup> irradiated autologous LPS (lipopolysaccharide)-blasts which had been pulsed with the respective peptide (100.0 µg per 10<sup>8</sup> cells). Cultures were kept in a 37°C, humidified CO<sub>2</sub> incubator for 6-7 days before being tested for effector function in a standard 4 hr *in vitro* <sup>51</sup>Cr-release CTL assay as follows. The responders were harvested from the 6-7 day cultures and washed twice with RPMI-1640 medium (without bovine serum). The positive target was created by incubating 3-5 x 10<sup>6</sup> P815-A2Kb transfectant with 100.0 µg of the specified peptide overnight in a 26°C water bath. The target cells were then labeled with <sup>51</sup>Cr at 250.0 µCi per 1 x 10<sup>6</sup> cells for 1 hr in the presence of 15.0 µg of the same test peptides. After washing twice with complete medium to remove excess free <sup>51</sup>Cr, the targets were incubated at 2.5 x 10<sup>3</sup> with different numbers of the responders for 4 hr in a 37°C CO<sub>2</sub> incubator. Supernatant aliquots were then removed and counted for radioactivity.

CLP-316 was selected as a representative PSA peptide for this study. The results depicted in Figure 1 reveal that CLP-316 was immunogenic and capable of eliciting an epitope-specific CTL response.

Whereas the invention is susceptible to various modifications and/or alternate forms, specific embodiments have been shown by way of example and are herein described in detail. However, it should be understood that it is not intended to limit the invention to the particular embodiments shown, but on the contrary, the invention is to cover all modifications, equivalents, and/or alternatives following within the spirit and scope of the invention as defined by the appended claims.

All publications, patents and patent applications referred to herein, are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.



Table 1. Capacity of PSA derived peptides to bind and stabilize HLA-A0201 molecules on T2 cells

Peptide	Amino Acid Position (Sequence)*	SEQ ID	Fluorescence Index (FI)
CLP-313	1-8 (MWVPVFL)	1	1.0
CLP-314	53-61 (VLVHPQWVL)	2	1.1
CLP-316	170-178 (KLQCVDLHV)	3	2.7
CLP-315	140-148 (DLPTQEPAL)	4	0
CLP-317	175-183 (DLHVISNDV)	5	0
CLP-320	99-107 (PLYDMSLLK)***	6	0.2**

\*Amino acid position pertains to that of the preproprotein.

\*\* Analysis did not include 3hr de-stabilization step

\*\*\* Negative Control

SEQUENCE LISTING

<110> Chong, Pele

Pedyczak, Artur

Sia, Charles Dwo Yuan

<120> Immunogenic Peptides Derived from Prostate-Specific Antigen (PSA) and  
Uses Thereof

<130> 11014-24

<150> US 60/195,456

<151> 2000-04-10

<160> 12

<170> PatentIn version 3.0

<210> 1

<211> 8

<212> PRT

<213> Artificial

<220>

<223> CLP313

<400> 1

Met Trp Val Pro Val Val Phe Leu  
1 5

<210> 2

<211> 9

<212> PRT

<213> Artificial

<220>

<223> CLP314

<400> 2

Val Leu Val His Pro Gln Trp Val Leu  
1 5

<210> 3

<211> 9

<212> PRT

<213> Artificial

<220>

<223> CLP316

<400> 3

Lys Leu Gln Cys Val Asp Leu His Val  
1 5

<210> 4

<211> 9

<212> PRT

<213> Artificial

<220>

<223> CLP315

<400> 4

Asp Leu Pro Thr Gln Glu Pro Ala Leu  
1 5

<210> 5

<211> 9

<212> PRT

<213> Artificial

<220>

<223> CLP317

<400> 5

Asp Leu His Val Ile Ser Asn Asp Val  
1 5

<210> 6

<211> 9

<212> PRT

<213> Artificial

<220>

<223> CLP320

<400> 6

Pro Leu Tyr Asp Met Ser Leu Leu Lys  
1 5

<210> 7  
<211> 24  
<212> DNA  
<213> Artificial

<220>  
<223> CLP313  
<400> 7  
atgtgggtcc cgggtgtctt cctc

24

<210> 8  
<211> 27  
<212> DNA  
<213> Artificial

<220>  
<223> CLP314  
<400> 8  
gttctgggtgc acccccagtg ggtctctc

27

<210> 9  
<211> 27  
<212> DNA  
<213> Artificial

<220>  
<223> CLP316  
<400> 9  
aaacttcagt gtgtggacct ccatgtt

27

<210> 10  
<211> 27  
<212> DNA  
<213> Artificial

<220>  
<223> CLP315  
<400> 10  
gacctgccca cccaggagcc agcactg

27

<210> 11

<211> 27

<212> DNA

<213> Artificial

<220>

<223> CLP317

<400> 11  
gacctccatg ttatttccaa tgacgtg

27

<210> 12

<211> 27

<212> DNA

<213> Artificial

<220>

<223> CLP320

<400> 12  
ccgctctacg atatgagcct cctgaag

27

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119